arXiv:1401.2946v1 [math.CA] 7 Jan 2014

ON GENERALIZATION OF DIFFERENT TYPE INEQUALITIES FOR HARMONICALLY QUASI-CONVEX FUNCTIONS VIA FRACTIONAL INTEGRALS

İMDAT İŞCAN

Abstract. In this paper, we obtained some new estimates on generalization of Hadamard, Ostrowski and Simpson-like type inequalities for harmonically quasi-convex functions via Riemann Liouville fractional integral.

1. Introduction

In this section we will present definitions and some results used in this paper.

A function $f : I \subseteq \mathbb{R} \to \mathbb{R}$ is said to be a convex function if

$$f(\lambda x + (1-\lambda)y) \leq \lambda f(x) + (1-\lambda)f(y)$$

holds for all $x, y \in I$ and $\lambda \in [0,1]$.

The notion of quasi-convex functions generalizes the notion of convex functions. More precisely, a function $f : [u,v] \to \mathbb{R}$ is said quasi-convex on $[u,v]$ if

$$f(\lambda x + (1-\lambda)y) \leq \sup\{f(x), f(y)\},$$

for any $x, y \in [u,v]$ and $\lambda \in [0,1]$. Clearly, any convex function is a quasi-convex function. Furthermore, there exist quasi-convex functions which are not convex (see [4]).

Following inequalities are well known in the literature as Hermite-Hadamard inequality, Ostrowski inequality and Simpson inequality respectively:

Theorem 1. *Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a convex function defined on the interval I of real numbers and $u, v \in I$ with $u < v$. The following double inequality holds*

$$f\left(\frac{u+v}{2}\right) \leq \frac{1}{v-u}\int_u^v f(x)dx \leq \frac{f(u)+f(v)}{2}. \tag{1.1}$$

Theorem 2. *Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a mapping differentiable in I°, the interior of I, and let $u, v \in I^\circ$ with $u < v$. If $|f'(x)| \leq M$ for all $x \in [u,v]$, then the following inequality holds*

$$\left| f(x) - \frac{1}{v-u}\int_u^v f(t)dt \right| \leq \frac{M}{v-u}\left[\frac{(x-u)^2 + (v-x)^2}{2}\right]$$

for all $x \in [u,v]$.

2000 *Mathematics Subject Classification.* 26A33, 26A51, 26D15.

Key words and phrases. Hermite–Hadamard inequality, Riemann–Liouville fractional integral, Ostrowski inequality, Simpson type inequalities, harmonically quasi-convex function.

Theorem 3. *Let $f:[u,v]\to\mathbb{R}$ be a four times continuously differentiable mapping on (u,v) and $\left\|f^{(4)}\right\|_\infty=\sup\limits_{x\in(u,v)}\left|f^{(4)}(x)\right|<\infty$. Then the following inequality holds:*

$$\left|\frac{1}{3}\left[\frac{f(u)+f(v)}{2}+2f\left(\frac{u+v}{2}\right)\right]-\frac{1}{v-u}\int_u^v f(x)dx\right|\leq\frac{1}{2880}\left\|f^{(4)}\right\|_\infty(v-u)^4.$$

In [5], Iscan defined harmonically convex functions and gave some Hermite-Hadamard type inequalities for this class of functions

Definition 1. *Let $I\subset\mathbb{R}\setminus\{0\}$ be a real interval. A function $f:I\to\mathbb{R}$ is said to be harmonically convex, if*

$$f\left(\frac{xy}{\lambda x+(1-\lambda)y}\right)\leq\lambda f(y)+(1-\lambda)f(x) \tag{1.2}$$

for all $x,y\in I$ and $\lambda\in[0,1]$. If the inequality in (1.2) is reversed, then f is said to be harmonically concave.

In [17], Zhang et al. defined the harmonically quasi-convex function and supplied several properties of this kind of functions.

Definition 2. *A function $f:I\subseteq(0,\infty)\to[0,\infty)$ is said to be harmonically convex, if*

$$f\left(\frac{xy}{\lambda x+(1-\lambda)y}\right)\leq\sup\{f(x),f(y)\}$$

for all $x,y\in I$ and $\lambda\in[0,1]$.

We would like to point out that any harmonically convex function on $I\subseteq(0,\infty)$ is a harmonically quasi-convex function, but not conversely. For example, the function

$$f(x)=\begin{cases}1, & x\in(0,1];\\ (x-2)^2, & x\in[1,4].\end{cases}$$

is harmonically quasi-convex on $(0,4]$, but it is not harmonically convex on $(0,4]$.

Let us recall the following special functions:

(1) The Beta function:

$$\beta(x,y)=\frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}=\int_0^1 t^{x-1}(1-t)^{y-1}dt,\quad x,y>0,$$

(2) The hypergeometric function:

$${}_2F_1(a,b;c;z)=\frac{1}{\beta(b,c-b)}\int_0^1 t^{b-1}(1-t)^{c-b-1}(1-zt)^{-a}dt,\ c>b>0,\ |z|<1\ \text{(see [11])}.$$

We give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used throughout this paper.

Definition 3. *Let $f\in L[u,v]$. The Riemann-Liouville integrals $J^\alpha_{u+}f$ and $J^\alpha_{v-}f$ of oder $\alpha>0$ with $u\geq0$ are defined by*

$$J_{u+}^{\alpha}f(b)=\frac{1}{\Gamma(\alpha)}\int_{u}^{b}(b-t)^{\alpha-1}f(t)dt,\ b>u$$

and

$$J_{v-}^{\alpha}f(a)=\frac{1}{\Gamma(\alpha)}\int_{a}^{v}(t-a)^{\alpha-1}f(t)dt,\ a<v$$

respectively, where $\Gamma(\alpha)$ *is the Gamma function defined by* $\Gamma(\alpha)=\int_{0}^{\infty}e^{-t}t^{\alpha-1}dt$ *and* $J_{u+}^{0}f(x)=J_{v-}^{0}f(x)=f(x)$.

In the case of $\alpha=1$, the fractional integral reduces to the classical integral. In recent years, many athors have studied errors estimations for Hermite-Hadamard, Ostrowski and Simpson inequalities; for refinements, counterparts, generalization see [1, 2, 3, 7, 8, 9, 10, 12, 13, 14, 15, 16].

In [6], Iscan gave the following new identity for differentiable functions and obtained some new general integral inequalities for harmonically convex functions via fractional integrals.

Lemma 1. *Let* $f:I\subseteq(0,\infty)\to\mathbb{R}$ *be a differentiable function on* I° *and* $a,b\in I$ *with* $a<b$. *If* $f'\in L[a,b]$ *then for all* $x\in[a,b]$, $\lambda\in[0,1]$ *and* $\alpha>0$ *we have:*

$$I_{f,g}\left(x,\lambda,\alpha,a,b\right)=\frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}}\int_{0}^{1}\frac{t^{\alpha}-\lambda}{A_{t}^{2}(a,x)}f'\left(\frac{ax}{A_{t}(a,x)}\right)dt$$

$$-\frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}}\int_{0}^{1}\frac{t^{\alpha}-\lambda}{A_{t}^{2}(b,x)}f'\left(\frac{bx}{A_{t}(b,x)}\right)dt,$$

where $A_{t}(u,x)=tu+(1-t)x$ *and*

$$\begin{aligned}(1.3)\quad &I_{f,g}\left(x,\lambda,\alpha,a,b\right)\\ =\ &(1-\lambda)\left[\left(\frac{x-a}{ax}\right)^{\alpha}+\left(\frac{b-x}{bx}\right)^{\alpha}\right]f(x)+\lambda\left[\left(\frac{x-a}{ax}\right)^{\alpha}f(a)+\left(\frac{b-x}{bx}\right)^{\alpha}f(b)\right]\\ &-\Gamma\left(\alpha+1\right)\left[J_{1/x+}^{\alpha}\left(f\circ g\right)\left(1/a\right)+J_{1/x-}^{\alpha}\left(f\circ g\right)\left(1/b\right)\right],\ g(u)=1/u.\end{aligned}$$

In this paper, by using of Lemma 1, we obtained a generalization of Hadamard, Ostrowski and Simpson type inequalities for harmonically quasi-convex functions via Riemann Liouville fractional integral.

2. Main Results

Let $f:I\subseteq(0,\infty)\to\mathbb{R}$ be a differentiable function on I°, the interior of I, throughout this section we will take the notation $I_{f,g}\left(x,\lambda,\alpha,a,b\right)$ as in (1.3), where $a,b\in I$ with $a<b$, $x\in[a,b]$, $\lambda\in[0,1]$, $g(u)=1/u$, $\alpha>0$ and Γ is Euler Gamma function. In order to prove our main results we need the following identity.

Theorem 4. *Let $f : I \subseteq (0,\infty) \to \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a,b]$, where $a,b \in I^\circ$ with $a < b$. If $|f'|^q$ is harmonically quasi-convex on $[a,b]$ for some fixed $q \geq 1$, then for $x \in [a,b]$, $\lambda \in [0,1]$ and $\alpha > 0$ the following inequality holds*

$$
\begin{aligned}
&|I_{f,g}(x,\lambda,\alpha,a,b)| \leq C_1^{1-1/q}(\alpha,\lambda) \\
&\times \left\{ \frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}x^{2q}} \left[\sup\left\{|f'(x)|^q, |f'(a)|^q\right\}\right]^{1/q} C_2^{1/q}\left(\alpha,\lambda,q,\frac{a}{x}\right) \right. \\
&\left. + \frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}b^{2q}} \left[\sup\left\{|f'(x)|^q, |f'(a)|^q\right\}\right]^{1/q} C_3^{1/q}\left(\alpha,\lambda,q,\frac{x}{b}\right) \right\},
\end{aligned} \tag{2.1}
$$

where

$$C_1(\alpha,\lambda) = \frac{2\alpha\lambda^{1+\frac{1}{\alpha}}+1}{\alpha+1} - \lambda,$$

$$
\begin{aligned}
&C_2\left(\alpha,\lambda,q,\frac{a}{x}\right) \\
= \;& \frac{1}{\alpha+1}\cdot{}_2F_1(2q;\alpha+1;\alpha+2,1-\frac{a}{x}) - \lambda\cdot{}_2F_1(2q;1;2,1-\frac{a}{x}) \\
& +2\lambda^{1+\frac{1}{\alpha}}\cdot\left[{}_2F_1(2q;1;2,\lambda^{1/\alpha}\left(1-\frac{a}{x}\right)) - \frac{1}{\alpha+1}\cdot{}_2F_1(2q;\alpha+1;\alpha+2,\lambda^{1/\alpha}\left(1-\frac{a}{x}\right))\right],
\end{aligned}
$$

$$
\begin{aligned}
&C_3\left(\alpha,\lambda,q,\frac{x}{b}\right) \\
= \;& \frac{1}{\alpha+1}\cdot{}_2F_1(2q;1;\alpha+2,1-\frac{x}{b}) - \lambda\cdot{}_2F_1(2q;1;2,1-\frac{x}{b}) \\
& +2\lambda^{1+\frac{1}{\alpha}}\cdot\left[{}_2F_1(2q;1;2,\lambda^{1/\alpha}\left(1-\frac{x}{b}\right)) - \frac{1}{\alpha+1}\cdot{}_2F_1(2q;1;\alpha+2,\lambda^{1/\alpha}\left(1-\frac{x}{b}\right))\right].
\end{aligned}
$$

Proof. Since$|f'|^q$ is harmonically quasi-convex on $[a,b]$, from Lemma 1, property of the modulus and using the power-mean inequality we have

$$
\begin{aligned}
&|I_{f,g}(x,\lambda,\alpha,a,b)| \leq \frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}} \int_0^1 \frac{|t^\alpha-\lambda|}{A_t^2(a,x)} \left|f'\left(\frac{ax}{A_t(a,x)}\right)\right| dt \\
&+\frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}} \int_0^1 \frac{|t^\alpha-\lambda|}{A_t^2(b,x)} \left|f'\left(\frac{bx}{A_t(b,x)}\right)\right| dt \\
\leq \;& \frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}} \left(\int_0^1 |t^\alpha-\lambda|\,dt\right)^{1-\frac{1}{q}} \left(\int_0^1 \frac{|t^\alpha-\lambda|}{A_t^{2q}(a,x)} \left|f'\left(\frac{ax}{A_t(a,x)}\right)\right|^q dt\right)^{\frac{1}{q}} \\
&+\frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}} \left(\int_0^1 |t^\alpha-\lambda|\,dt\right)^{1-\frac{1}{q}} \left(\int_0^1 \frac{|t^\alpha-\lambda|}{A_t^{2q}(b,x)} \left|f'\left(\frac{bx}{A_t(b,x)}\right)\right|^q dt\right)^{\frac{1}{q}}
\end{aligned}
$$

$$\leq C_1^{1-1/q}(\alpha,\lambda)\left\{\frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}}\left[\sup\left\{|f'(x)|^q,|f'(a)|^q\right\}\right]^{1/q}\left(\int_0^1\frac{|t^\alpha-\lambda|}{A_t^{2q}(a,x)}dt\right)^{1/q}\right.$$

$$\left.+\frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}}\left[\sup\left\{|f'(x)|^q,|f'(a)|^q\right\}\right]^{1/q}\left(\int_0^1\frac{|t^\alpha-\lambda|}{A_t^{2q}(a,x)}dt\right)^{1/q}\right\} \tag{2.2}$$

where by simple computation we obtain

$$C_1(\alpha,\lambda) = \int_0^1|t^\alpha-\lambda|\,dt$$

$$= \frac{2\alpha\lambda^{1+\frac{1}{\alpha}}+1}{\alpha+1}-\lambda, \tag{2.3}$$

$$\int_0^1\frac{|t^\alpha-\lambda|}{A_t^{2q}(a,x)}dt \tag{2.4}$$

$$= x^{-2q}\left\{\frac{1}{\alpha+1}\cdot{}_2F_1(2q;\alpha+1;\alpha+2,1-\frac{a}{x})-\lambda\cdot{}_2F_1(2q;1;2,1-\frac{a}{x})\right.$$

$$\left.+2\lambda^{1+\frac{1}{\alpha}}\cdot\left[{}_2F_1(2q;1;2,\lambda^{1/\alpha}\left(1-\frac{a}{x}\right))-\frac{1}{\alpha+1}\cdot{}_2F_1(2q;\alpha+1;\alpha+2,\lambda^{1/\alpha}\left(1-\frac{a}{x}\right))\right],$$

and

$$\int_0^1\frac{|t^\alpha-\lambda|}{A_t^{2q}(b,x)}dt \tag{2.5}$$

$$= b^{-2q}\left\{\frac{1}{\alpha+1}\cdot{}_2F_1(2q;1;\alpha+2,1-\frac{x}{b})-\lambda\cdot{}_2F_1(2q;1;2,1-\frac{x}{b})\right.$$

$$\left.+2\lambda^{1+\frac{1}{\alpha}}\cdot\left[{}_2F_1(2q;1;2,\lambda^{1/\alpha}\left(1-\frac{x}{b}\right))-\frac{1}{\alpha+1}\cdot{}_2F_1(2q;1;\alpha+2,\lambda^{1/\alpha}\left(1-\frac{x}{b}\right))\right],$$

Hence, If we use (2.3), (2.4) and (2.5) in (2.2), we obtain the desired result. This completes the proof. □

Corollary 1. *Under the assumptions of Theorem 4 with $q=1$, the inequality (2.1) reduced to the following inequality*

$$|I_{f,g}(x,\lambda,\alpha,a,b)|$$

$$\leq \left\{\frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}x^2}\left[\sup\left\{|f'(x)|,|f'(a)|\right\}\right]C_2\left(\alpha,\lambda,1,\frac{a}{x}\right)\right.$$

$$\left.+\frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}b^2}\left[\sup\left\{|f'(x)|,|f'(a)|\right\}\right]^{1/q}C_3\left(\alpha,\lambda,1,\frac{x}{b}\right)\right\},$$

Corollary 2. *Under the assumptions of Theorem 4 with $\alpha = 1$, the inequality (2.1) reduced to the following inequality*

$$\begin{aligned}
&\left(\frac{ab}{b-a}\right)\left|I_{f,g}\left(x,\lambda,\alpha,a,b\right)\right| \\
= &\left|\left(1-\lambda\right)f\left(x\right)+\lambda\left[\frac{b(x-a)f(a)+a(b-x)f(b)}{x(b-a)}\right]-\frac{ab}{b-a}\int_a^b\frac{f\left(u\right)}{u^2}du\right| \\
\leq &\left(\frac{ab}{b-a}\right)\left(\frac{2\lambda^2-2\lambda+1}{2}\right)^{1-\frac{1}{q}}\left\{\frac{\left(x-a\right)^2}{x^{2q}}\left[\sup\left\{\left|f'\left(x\right)\right|^q,\left|f'\left(a\right)\right|^q\right\}\right]^{1/q}C_2^{1/q}\left(1,\lambda,q,\frac{a}{x}\right)\right. \\
&\left.+\frac{\left(b-x\right)^2}{b^{2q}}\left[\sup\left\{\left|f'\left(x\right)\right|^q,\left|f'\left(a\right)\right|^q\right\}\right]^{1/q}C_3^{1/q}\left(1,\lambda,q,\frac{x}{b}\right)\right\},
\end{aligned}$$

specially for $x = H = 2ab/(a+b)$, we get

$$\begin{aligned}
&\left|\left(1-\lambda\right)f\left(H\right)+\lambda\left(\frac{f(a)+f(b)}{2}\right)-\frac{ab}{b-a}\int_a^b\frac{f\left(u\right)}{u^2}du\right|\leq\frac{b-a}{4ab}\left(\frac{2\lambda^2-2\lambda+1}{2}\right)^{1-\frac{1}{q}} \\
&\times\left\{\frac{a^2H^2}{H^{2q}}C_2^{1/q}\left(1,\lambda,q,\frac{a+b}{2b}\right)\left(\sup\left\{\left|f'\left(H\right)\right|^q,\left|f'\left(a\right)\right|^q\right\}\right)^{\frac{1}{q}}\right. \\
&\left.+\frac{b^2H^2}{b^{2q}}C_3^{1/q}\left(1,\lambda,q,\frac{2a}{a+b}\right)\left(\sup\left\{\left|f'\left(H\right)\right|^q,\left|f'\left(b\right)\right|^q\right\}\right)^{\frac{1}{q}}\right\}.
\end{aligned}$$

Corollary 3. *In Theorem 4,*

(1) If we take $x = H = 2ab/(a+b)$, $\lambda = \frac{1}{3}$, then we get the following Simpson type inequality for fractional integrals

$$\begin{aligned}
&\left|\frac{1}{6}\left[f(a)+4f\left(H\right)+f(b)\right]-\left(\frac{ab}{b-a}\right)^{\alpha}2^{\alpha-1}\Gamma\left(\alpha+1\right)\left[J_{1/H+}^{\alpha}\left(f\circ g\right)\left(1/a\right)+J_{1/H-}^{\alpha}\left(f\circ g\right)\left(1/b\right)\right]\right| \\
\leq &\frac{b-a}{4ab}C_1^{1-1/q}\left(\alpha,\frac{1}{3}\right)\left\{\frac{a^2H^2}{H^{2q}}\left[\sup\left\{\left|f'\left(H\right)\right|^q,\left|f'\left(a\right)\right|^q\right\}\right]^{1/q}C_2^{1/q}\left(\alpha,\frac{1}{3},q,\frac{a}{H}\right)\right. \\
&\left.+\frac{b^2H^2}{b^{2q}}\left[\sup\left\{\left|f'\left(H\right)\right|^q,\left|f'\left(a\right)\right|^q\right\}\right]^{1/q}C_3^{1/q}\left(\alpha,\frac{1}{3},q,\frac{H}{b}\right)\right\},
\end{aligned}$$

specially for $\alpha = 1$, we get

$$\begin{aligned}
&\left|\frac{1}{6}\left[f(a)+4f\left(H\right)+f(b)\right]-\frac{ab}{b-a}\int_a^b\frac{f(u)}{u^2}du\right|\leq\frac{b-a}{4ab}\left(\frac{5}{18}\right)^{1-\frac{1}{q}} \\
&\times\left\{\frac{a^2H^2}{H^{2q}}C_2^{1/q}\left(1,\frac{1}{3},q,\frac{a+b}{2b}\right)\left[\sup\left\{\left|f'\left(H\right)\right|,\left|f'\left(a\right)\right|\right\}\right]^{\frac{1}{q}}\right. \\
&\left.+\frac{b^2H^2}{b^{2q}}C_3^{1/q}\left(1,\frac{1}{3},q,\frac{2a}{a+b}\right)\left[\sup\left\{\left|f'\left(H\right)\right|^q,\left|f'\left(b\right)\right|^q\right\}\right]^{\frac{1}{q}}\right\}.
\end{aligned}$$

(2) If we take $x = H = 2ab/(a+b)$, $\lambda = 0$, then we get the following midpoint type inequality for fractional integrals

$$\begin{aligned}
&\left| f(H) - \left(\frac{ab}{b-a}\right)^{\alpha} 2^{\alpha-1}\Gamma(\alpha+1)\left[J^{\alpha}_{1/H+}(f\circ g)(1/a) + J^{\alpha}_{1/H-}(f\circ g)(1/b)\right]\right| \\
\leq\ & \frac{b-a}{4ab}\left(\frac{1}{\alpha+1}\right)^{1-\frac{1}{q}}\left\{\frac{a^2H^2}{H^{2q}}C_2^{1/q}\left(\alpha,0,q,\frac{a+b}{2b}\right)\left[\sup\left\{|f'(H)|^q,|f'(a)|^q\right\}\right]^{\frac{1}{q}}\right. \\
&\left.+\frac{b^2H^2}{b^{2q}}C_3^{1/q}\left(\alpha,0,q,\frac{2a}{a+b}\right)\left[\sup\left\{|f'(H)|^q,|f'(b)|^q\right\}\right]^{\frac{1}{q}}\right\},
\end{aligned}$$

specially for $\alpha = 1$, we get

$$\begin{aligned}
&\left| f(H) - \frac{ab}{b-a}\int_a^b \frac{f(u)}{u^2}du\right| \\
\leq\ & \frac{b-a}{4ab}\left(\frac{1}{2}\right)^{1-\frac{1}{q}}\left\{\frac{a^2H^2}{H^{2q}}C_2^{1/q}\left(1,0,q,\frac{a+b}{2b}\right)\left[\sup\left\{|f'(H)|^q,|f'(a)|^q\right\}\right]^{\frac{1}{q}}\right. \\
&\left.+\frac{b^2H^2}{b^{2q}}C_3^{1/q}\left(1,0,q,\frac{2a}{a+b}\right)\left[\sup\left\{|f'(H)|^q,|f'(b)|^q\right\}\right]^{\frac{1}{q}}\right\}.
\end{aligned}$$

(3) If we take $x = H = 2ab/(a+b)$, $\lambda = 1$, then we get the following trapezoid type inequality for fractional integrals

$$\begin{aligned}
&\left| \frac{f(a)+f(b)}{2} - \left(\frac{ab}{b-a}\right)^{\alpha} 2^{\alpha-1}\Gamma(\alpha+1)\left[J^{\alpha}_{1/H+}(f\circ g)(1/a) + J^{\alpha}_{1/H-}(f\circ g)(1/b)\right]\right| \\
\leq\ & \frac{b-a}{4ab}\left(\frac{\alpha}{\alpha+1}\right)^{1-\frac{1}{q}}\left\{\frac{a^2H^2}{H^{2q}}C_2^{1/q}\left(\alpha,1,q,\frac{a+b}{2b}\right)\left[\sup\left\{|f'(H)|^q,|f'(a)|^q\right\}\right]^{\frac{1}{q}}\right. \\
&\left.+\frac{b^2H^2}{b^{2q}}C_3^{1/q}\left(\alpha,1,q,\frac{2a}{a+b}\right)\left[\sup\left\{|f'(H)|^q,|f'(b)|^q\right\}\right]^{\frac{1}{q}}\right\},
\end{aligned}$$

specially for $\alpha = 1$, we get

$$\begin{aligned}
&\left| \frac{f(a)+f(b)}{2} - \frac{ab}{b-a}\int_a^b \frac{f(u)}{u^2}du\right| \\
\leq\ & \frac{b-a}{4ab}\left(\frac{1}{2}\right)^{1-\frac{1}{q}}\left\{\frac{a^2H^2}{H^{2q}}C_2^{1/q}\left(1,1,q,\frac{a+b}{2b}\right)\left[\sup\left\{|f'(H)|^q,|f'(a)|^q\right\}\right]^{\frac{1}{q}}\right. \\
&\left.+\frac{b^2H^2}{b^{2q}}C_3^{1/q}\left(1,1,q,\frac{2a}{a+b}\right)\left[\sup\left\{|f'(H)|^q,|f'(b)|^q\right\}\right]^{\frac{1}{q}}\right\}.
\end{aligned}$$

Corollary 4. *Let the assumptions of Theorem 4 hold. If $|f'(x)| \leq M$ for all $x \in [a,b]$ and $\lambda = 0$, then we get the following Ostrowski type inequality for fractional from the inequality (2.1) integrals*

$$\begin{aligned}
&\left|\left[\left(\frac{x-a}{ax}\right)^{\alpha} + \left(\frac{b-x}{bx}\right)^{\alpha}\right]f(x) - \left(\frac{ab}{b-a}\right)^{\alpha} 2^{\alpha-1}\Gamma(\alpha+1)\left[J^{\alpha}_{1/x+}(f\circ g)(1/a) + J^{\alpha}_{1/x-}(f\circ g)(1/b)\right]\right| \\
\leq\ & \frac{M}{(\alpha+1)^{1-\frac{1}{q}}}\left[\frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}x^{2q}}C_2^{1/q}\left(\alpha,0,q,\frac{a}{x}\right) + \frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}b^{2q}}C_3^{1/q}\left(\alpha,0,q,\frac{x}{b}\right)\right].
\end{aligned}$$

Theorem 5. *Let $f : I \subseteq (0,\infty) \to \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a,b]$, where $a,b \in I^\circ$ with $a < b$. If $|f'|^q$ is harmonically quasi-convex on $[a,b]$ for some fixed $q \geq 1$, then for $x \in [a,b]$, $\lambda \in [0,1]$ and $\alpha > 0$ the following inequality holds*

$$\begin{aligned}(2.6)\quad |I_{f,g}(x,\lambda,\alpha,a,b)| &\leq C_2\left(\alpha,\lambda,1,\frac{a}{x}\right)\frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}x^{2q}}\left[\sup\left\{|f'(x)|^q,|f'(a)|^q\right\}\right]^{1/q}\\ &\quad +C_3\left(\alpha,\lambda,1,\frac{x}{b}\right)\frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}b^{2q}}\left[\sup\left\{|f'(x)|^q,|f'(b)|^q\right\}\right]^{1/q}\end{aligned}$$

where C_2 and C_3 are defined as in Theorem 4.

Proof. Since$|f'|^q$ is harmonically quasi-convex on $[a,b]$, from Lemma 1,using property of the modulus and the power-mean inequality we have

$$\begin{aligned}&|I_{f,g}(x,\lambda,\alpha,a,b)|\\ &\leq \frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}}\left(\int_0^1\frac{|t^\alpha-\lambda|}{A_t^2(a,x)}dt\right)^{1-\frac{1}{q}}\left(\int_0^1\frac{|t^\alpha-\lambda|}{A_t^2(a,x)}\left|f'\left(\frac{ax}{A_t(a,x)}\right)\right|^q dt\right)^{\frac{1}{q}}\\ &\quad +\frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}}\left(\int_0^1\frac{|t^\alpha-\lambda|}{A_t^2(b,x)}dt\right)^{1-\frac{1}{q}}\left(\int_0^1\frac{|t^\alpha-\lambda|}{A_t^2(b,x)}\left|f'\left(\frac{bx}{A_t(b,x)}\right)\right|^q dt\right)^{\frac{1}{q}}\end{aligned}$$

$$\begin{aligned}&\leq C_2\left(\alpha,\lambda,1,\frac{a}{x}\right)\frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}x^{2q}}\left[\sup\left\{|f'(x)|^q,|f'(a)|^q\right\}\right]^{1/q}\\ &\quad +C_3\left(\alpha,\lambda,1,\frac{x}{b}\right)\frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}b^{2q}}\left[\sup\left\{|f'(x)|^q,|f'(b)|^q\right\}\right]^{1/q}\end{aligned}$$

which completes the proof. □

Corollary 5. *Under the assumptions of Theorem 5 with $q = 1$, the inequality (2.6) reduced to the following inequality*

$$\begin{aligned}|I_{f,g}(x,\lambda,\alpha,a,b)| &\leq C_2\left(\alpha,\lambda,1,\frac{a}{x}\right)\frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}x^2}\left[\sup\left\{|f'(x)|,|f'(a)|\right\}\right]\\ &\quad +C_3\left(\alpha,\lambda,1,\frac{x}{b}\right)\frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}b^2}\left[\sup\left\{|f'(x)|,|f'(b)|\right\}\right]\end{aligned}$$

Corollary 6. *Under the assumptions of Theorem 5 with $\alpha = 1$, the inequality (2.6) reduced to the following inequality*

$$\begin{aligned}
&\left(\frac{ab}{b-a}\right)|I_{f,g}(x,\lambda,\alpha,a,b)| \\
= &\left|(1-\lambda)f(x)+\lambda\left[\frac{b(x-a)f(a)+a(b-x)f(b)}{x(b-a)}\right]-\frac{ab}{b-a}\int_a^b\frac{f(u)}{u^2}du\right| \\
\leq &\left(\frac{ab}{b-a}\right)\left\{\frac{(x-a)^2}{x^2}\left[\sup\left\{|f'(x)|^q,|f'(a)|^q\right\}\right]^{1/q}C_2\left(1,\lambda,1,\frac{a}{x}\right)\right. \\
&\left.+\frac{(b-x)^2}{b^2}\left[\sup\left\{|f'(x)|^q,|f'(a)|^q\right\}\right]^{1/q}C_3\left(1,\lambda,1,\frac{x}{b}\right)\right\},
\end{aligned}$$

specially for $x = H = 2ab/(a+b)$, we get

$$\begin{aligned}
&\left|(1-\lambda)f(H)+\lambda\left(\frac{f(a)+f(b)}{2}\right)-\frac{ab}{b-a}\int_a^b\frac{f(u)}{u^2}du\right|\leq\frac{b-a}{4ab} \\
&\times\left\{a^2C_2\left(1,\lambda,1,\frac{a+b}{2b}\right)\left(\sup\left\{|f'(H)|^q,|f'(a)|^q\right\}\right)^{\frac{1}{q}}\right. \\
&\left.+H^2C_3\left(1,\lambda,1,\frac{2a}{a+b}\right)\left(\sup\left\{|f'(H)|^q,|f'(b)|^q\right\}\right)^{\frac{1}{q}}\right\}.
\end{aligned}$$

Corollary 7. *In Theorem 5,*

(1) If we take $x = H = 2ab/(a+b)$, $\lambda = \frac{1}{3}$, then we get the following Simpson type inequality for fractional integrals

$$\begin{aligned}
&\left|\frac{1}{6}[f(a)+4f(H)+f(b)]-\left(\frac{ab}{b-a}\right)^{\alpha}2^{\alpha-1}\Gamma(\alpha+1)\left[J^{\alpha}_{1/H+}(f\circ g)(1/a)+J^{\alpha}_{1/H-}(f\circ g)(1/b)\right]\right| \\
\leq &\frac{b-a}{4ab}\left\{a^2\left[\sup\left\{|f'(H)|^q,|f'(a)|^q\right\}\right]^{1/q}C_2\left(\alpha,\frac{1}{3},1,\frac{a}{x}\right)\right. \\
&\left.+H^2\left[\sup\left\{|f'(H)|^q,|f'(a)|^q\right\}\right]^{1/q}C_3\left(\alpha,\frac{1}{3},1,\frac{x}{b}\right)\right\},
\end{aligned}$$

specially for $\alpha = 1$, we get

$$\begin{aligned}
&\left|\frac{1}{6}[f(a)+4f(H)+f(b)]-\frac{ab}{b-a}\int_a^b\frac{f(u)}{u^2}du\right|\leq\frac{b-a}{4ab} \\
&\times\left\{a^2C_2\left(1,\frac{1}{3},1,\frac{a+b}{2b}\right)\left[\sup\left\{|f'(H)|,|f'(a)|\right\}\right]^{\frac{1}{q}}\right. \\
&\left.+H^2C_3\left(1,\frac{1}{3},1,\frac{2a}{a+b}\right)\left[\sup\left\{|f'(H)|^q,|f'(b)|^q\right\}\right]^{\frac{1}{q}}\right\}.
\end{aligned}$$

(2) If we take $x=H=2ab/(a+b)$, $\lambda=0$, then we get the following midpoint type inequality for fractional integrals

$$\begin{aligned}&\left|f(H)-\left(\frac{ab}{b-a}\right)^{\alpha}2^{\alpha-1}\Gamma(\alpha+1)\left[J^{\alpha}_{1/H+}(f\circ g)(1/a)+J^{\alpha}_{1/H-}(f\circ g)(1/b)\right]\right|\\ \leq\ &\frac{b-a}{4ab}\left\{a^2C_2\left(\alpha,0,1,\frac{a+b}{2b}\right)\left[\sup\left\{|f'(H)|^q,|f'(a)|^q\right\}\right]^{\frac{1}{q}}\right.\\ &\left.+H^2C_3\left(\alpha,0,1,\frac{2a}{a+b}\right)\left[\sup\left\{|f'(H)|^q,|f'(b)|^q\right\}\right]^{\frac{1}{q}}\right\},\end{aligned}$$

specially for $\alpha=1$, we get

$$\begin{aligned}&\left|f(H)-\frac{ab}{b-a}\int_a^b\frac{f(u)}{u^2}du\right|\\ \leq\ &\frac{b-a}{4ab}\left(\frac{1}{2}\right)^{1-\frac{1}{q}}\left\{\frac{a^2H^2}{H^{2q}}C_2^{1/q}\left(1,0,q,\frac{a+b}{2b}\right)\left[\sup\left\{\left|f'\left(\frac{2ab}{a+b}\right)\right|^q,|f'(a)|^q\right\}\right]^{\frac{1}{q}}\right.\\ &\left.+\frac{b^2H^2}{b^{2q}}C_3^{1/q}\left(1,0,q,\frac{2a}{a+b}\right)\left[\sup\left\{\left|f'\left(\frac{2ab}{a+b}\right)\right|^q,|f'(b)|^q\right\}\right]^{\frac{1}{q}}\right\}.\end{aligned}$$

(3) If we take $x=H=2ab/(a+b)$, $\lambda=1$, then we get the following trapezoid type inequality for fractional integrals

$$\begin{aligned}&\left|\frac{f(a)+f(b)}{2}-\left(\frac{ab}{b-a}\right)^{\alpha}2^{\alpha-1}\Gamma(\alpha+1)\left[J^{\alpha}_{1/H+}(f\circ g)(1/a)+J^{\alpha}_{1/H-}(f\circ g)(1/b)\right]\right|\\ \leq\ &\frac{b-a}{4ab}\left\{a^2C_2\left(\alpha,1,1,\frac{a+b}{2b}\right)\left[\sup\left\{|f'(H)|^q,|f'(a)|^q\right\}\right]^{\frac{1}{q}}\right.\\ &\left.+H^2C_3\left(\alpha,1,1,\frac{2a}{a+b}\right)\left[\sup\left\{|f'(H)|^q,|f'(b)|^q\right\}\right]^{\frac{1}{q}}\right\},\end{aligned}$$

specially for $\alpha=1$, we get

$$\begin{aligned}&\left|\frac{f(a)+f(b)}{2}-\frac{ab}{b-a}\int_a^b\frac{f(u)}{u^2}du\right|\\ \leq\ &\frac{b-a}{4ab}\left\{a^2C_2^{1/q}\left(1,1,1,\frac{a+b}{2b}\right)\left[\sup\left\{|f'(H)|^q,|f'(a)|^q\right\}\right]^{\frac{1}{q}}\right.\\ &\left.+H^2C_3^{1/q}\left(1,1,1,\frac{2a}{a+b}\right)\left[\sup\left\{|f'(H)|^q,|f'(b)|^q\right\}\right]^{\frac{1}{q}}\right\}.\end{aligned}$$

Corollary 8. *Let the assumptions of Theorem 5 hold. If $|f'(x)|\leq M$ for all $x\in[a,b]$ and $\lambda=0$, then we get the following Ostrowski type inequality for fractional from the inequality (2.6) integrals*

$$\begin{aligned}&\left|\left[\left(\frac{x-a}{ax}\right)^{\alpha}+\left(\frac{b-x}{bx}\right)^{\alpha}\right]f(x)-\left(\frac{ab}{b-a}\right)^{\alpha}2^{\alpha-1}\Gamma(\alpha+1)\left[J^{\alpha}_{1/x+}(f\circ g)(1/a)+J^{\alpha}_{1/x-}(f\circ g)(1/b)\right]\right|\\ \leq\ &M\left[\frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}x^2}C_2\left(\alpha,0,1,\frac{a}{x}\right)+\frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}b^2}C_3\left(\alpha,0,1,\frac{x}{b}\right)\right].\end{aligned}$$

Theorem 6. *Let $f : I \subseteq (0,\infty) \to \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a,b]$, where $a,b \in I^\circ$ with $a < b$. If $|f'|^q$ is harmonically quasi-convex on $[a,b]$ for some fixed $q > 1$, then for $x \in [a,b]$, $\lambda \in [0,1]$ and $\alpha > 0$ the following inequality holds*

$$
\begin{aligned}
(2.7)\qquad & |I_{f,g}(x,\lambda,\alpha,a,b)| \\
\leq\ & C_1^{1/q}(\alpha,\lambda)\left\{C_2^{1/p}\left(\alpha,\lambda,p,\frac{a}{x}\right)\frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}x^{2p}}\left[\sup\left\{|f'(x)|^q,|f'(a)|^q\right\}\right]^{1/q}\right. \\
& \left.+C_3^{1/p}\left(\alpha,\lambda,p,\frac{x}{b}\right)\frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}b^{2p}}\left[\sup\left\{|f'(x)|^q,|f'(b)|^q\right\}\right]^{1/q}\right\}
\end{aligned}
$$

where C_1, C_2 and C_3 are defined as in Theorem 4.

Proof. Since$|f'|^q$ is harmonically quasi-convex on $[a,b]$, from Lemma 1, using property of the modulus and the Hölder inequality we have

$$
\begin{aligned}
& |S_f(x,\lambda,\alpha,a,b)| \\
\leq\ & \frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}}\left(\int_0^1\frac{|t^\alpha-\lambda|}{A_t^{2p}(a,x)}dt\right)^{\frac{1}{p}}\left(\int_0^1|t^\alpha-\lambda|\left|f'\left(\frac{ax}{A_t(a,x)}\right)\right|^q dt\right)^{\frac{1}{q}} \\
& +\frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}}\left(\int_0^1\frac{|t^\alpha-\lambda|}{A_t^{2p}(b,x)}dt\right)^{\frac{1}{p}}\left(\int_0^1|t^\alpha-\lambda|\left|f'\left(\frac{bx}{A_t(b,x)}\right)\right|^q dt\right)^{\frac{1}{q}}
\end{aligned}
$$

$$
\begin{aligned}
\leq\ & C_1^{1/q}(\alpha,\lambda)\left\{C_2^{1/p}\left(\alpha,\lambda,p,\frac{a}{x}\right)\frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}x^{2p}}\left[\sup\left\{|f'(x)|^q,|f'(a)|^q\right\}\right]^{1/q}\right. \\
& \left.+C_3^{1/p}\left(\alpha,\lambda,p,\frac{x}{b}\right)\frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}b^{2p}}\left[\sup\left\{|f'(x)|^q,|f'(b)|^q\right\}\right]^{1/q}\right\}
\end{aligned}
$$

which completes the proof. □

Corollary 9. *Under the assumptions of Theorem 6 with $\alpha = 1$, the inequality (2.7) reduced to the following inequality*

$$
\begin{aligned}
& \left(\frac{ab}{b-a}\right)|I_{f,g}(x,\lambda,\alpha,a,b)| \\
=\ & \left|(1-\lambda)f(x)+\lambda\left[\frac{b(x-a)f(a)+a(b-x)f(b)}{x(b-a)}\right]-\frac{ab}{b-a}\int_a^b\frac{f(u)}{u^2}du\right| \\
\leq\ & \left(\frac{ab}{b-a}\right)\left(\frac{2\lambda^2-2\lambda+1}{2}\right)^{\frac{1}{q}}\left\{\frac{(x-a)^2}{x^{2p}}\left[\sup\left\{|f'(x)|^q,|f'(a)|^q\right\}\right]^{1/q}C_2^{1/p}\left(1,\lambda,p,\frac{a}{x}\right)\right. \\
& \left.+\frac{(b-x)^2}{b^{2p}}\left[\sup\left\{|f'(x)|^q,|f'(a)|^q\right\}\right]^{1/q}C_3^{1/p}\left(1,\lambda,p,\frac{x}{b}\right)\right\},
\end{aligned}
$$

specially for $x=H=2ab/(a+b)$, we get

$$\left|(1-\lambda)f(H)+\lambda\left(\frac{f(a)+f(b)}{2}\right)-\frac{ab}{b-a}\int_a^b\frac{f(u)}{u^2}du\right|\leq\frac{b-a}{4ab}\left(\frac{2\lambda^2-2\lambda+1}{2}\right)^{\frac{1}{q}}$$
$$\times\left\{\frac{a^2H^2}{H^{2p}}C_2^{1/p}\left(1,\lambda,p,\frac{a+b}{2b}\right)\left(\sup\left\{\left|f'(H)\right|^q,\left|f'(a)\right|^q\right\}\right)^{\frac{1}{q}}\right.$$
$$\left.+\frac{b^2H^2}{b^{2p}}C_3^{1/p}\left(1,\lambda,p,\frac{2a}{a+b}\right)\left(\sup\left\{\left|f'(H)\right|^q,\left|f'(b)\right|^q\right\}\right)^{\frac{1}{q}}\right\}.$$

Corollary 10. *In Theorem 6,*

(1) If we take $x=H=2ab/(a+b)$, $\lambda=\frac{1}{3}$, then we get the following Simpson type inequality for fractional integrals

$$\left(\frac{ab}{b-a}\right)^{\alpha}\left|I_{f,g}(x,\lambda,\alpha,a,b)\right|$$
$$\leq\frac{b-a}{4ab}C_1^{1-1/q}\left(\alpha,\frac{1}{3}\right)\left\{\frac{a^2H^2}{H^{2p}}\left[\sup\left\{\left|f'(H)\right|^q,\left|f'(a)\right|^q\right\}\right]^{1/q}C_2^{1/p}\left(\alpha,\frac{1}{3},p,\frac{a+b}{2b}\right)\right.$$
$$\left.+\frac{b^2H^2}{b^{2p}}\left[\sup\left\{\left|f'(H)\right|^q,\left|f'(a)\right|^q\right\}\right]^{1/q}C_3^{1/p}\left(\alpha,\frac{1}{3},p,\frac{2a}{a+b}\right)\right\},$$

specially for $\alpha=1$, we get

$$\left|\frac{1}{6}\left[f(a)+4f(H)+f(b)\right]-\frac{ab}{b-a}\int_a^b\frac{f(u)}{u^2}du\right|\leq\frac{b-a}{4ab}\left(\frac{5}{18}\right)^{\frac{1}{q}}$$
$$\times\left\{\frac{a^2H^2}{H^{2p}}C_2^{1/p}\left(1,\frac{1}{3},p,\frac{a+b}{2b}\right)\left[\sup\left\{\left|f'(H)\right|,\left|f'(a)\right|\right\}\right]^{\frac{1}{q}}\right.$$
$$\left.+\frac{b^2H^2}{b^{2p}}C_3^{1/p}\left(1,\frac{1}{3},p,\frac{2a}{a+b}\right)\left[\sup\left\{\left|f'(H)\right|^q,\left|f'(b)\right|^q\right\}\right]^{\frac{1}{q}}\right\}.$$

(2) If we take $x=H=2ab/(a+b)$, $\lambda=0$, then we get the following midpoint type inequality for fractional integrals

$$\left|f(H)-\left(\frac{ab}{b-a}\right)^{\alpha}2^{\alpha-1}\Gamma(\alpha+1)\left[J^{\alpha}_{1/H+}(f\circ g)(1/a)+J^{\alpha}_{1/H-}(f\circ g)(1/b)\right]\right|$$
$$\leq\frac{b-a}{4ab}\left(\frac{1}{\alpha+1}\right)^{\frac{1}{q}}\left\{\frac{a^2H^2}{H^{2p}}C_2^{1/p}\left(\alpha,0,p,\frac{a+b}{2b}\right)\left[\sup\left\{\left|f'(H)\right|^q,\left|f'(a)\right|^q\right\}\right]^{\frac{1}{q}}\right.$$
$$\left.+\frac{b^2H^2}{b^{2p}}C_3^{1/p}\left(\alpha,0,p,\frac{2a}{a+b}\right)\left[\sup\left\{\left|f'(H)\right|^q,\left|f'(b)\right|^q\right\}\right]^{\frac{1}{q}}\right\},$$

specially for $\alpha=1$, we get

$$\left|f(H)-\frac{ab}{b-a}\int_a^b\frac{f(u)}{u^2}du\right|$$
$$\leq\frac{b-a}{4ab}\left(\frac{1}{2}\right)^{\frac{1}{q}}\left\{\frac{a^2H^2}{H^{2p}}C_2^{1/p}\left(1,0,p,\frac{a+b}{2b}\right)\left[\sup\left\{\left|f'(H)\right|^q,\left|f'(a)\right|^q\right\}\right]^{\frac{1}{q}}\right.$$
$$\left.+\frac{b^2H^2}{b^{2p}}C_3^{1/p}\left(1,0,p,\frac{2a}{a+b}\right)\left[\sup\left\{\left|f'(H)\right|^q,\left|f'(b)\right|^q\right\}\right]^{\frac{1}{q}}\right\}.$$

(3) If we take $x = H = 2ab/(a+b)$, $\lambda = 1$, then we get the following trapezoid type inequality for fractional integrals

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2} - \left(\frac{ab}{b-a}\right)^{\alpha} 2^{\alpha-1}\Gamma(\alpha+1)\left[J^{\alpha}_{1/H+}(f\circ g)(1/a) + J^{\alpha}_{1/H-}(f\circ g)(1/b)\right]\right| \\
\leq\ & \frac{b-a}{4ab}\left(\frac{\alpha}{\alpha+1}\right)^{\frac{1}{q}}\left\{\frac{a^2H^2}{H^{2p}}C_2^{1/p}\left(\alpha,1,p,\frac{a+b}{2b}\right)\left[\sup\left\{|f'(H)|^q,|f'(a)|^q\right\}\right]^{\frac{1}{q}}\right. \\
&\left.+\frac{b^2H^2}{b^{2q}}C_3^{1/p}\left(\alpha,1,p,\frac{2a}{a+b}\right)\left[\sup\left\{|f'(H)|^q,|f'(b)|^q\right\}\right]^{\frac{1}{q}}\right\},
\end{aligned}$$

specially for $\alpha = 1$, we get

$$\begin{aligned}
&\left|\frac{f(a)+f(b)}{2} - \frac{ab}{b-a}\int_a^b \frac{f(u)}{u^2}du\right| \\
\leq\ & \frac{b-a}{4ab}\left(\frac{1}{2}\right)^{\frac{1}{q}}\left\{\frac{a^2H^2}{H^{2p}}C_2^{1/p}\left(1,1,p,\frac{a+b}{2b}\right)\left[\sup\left\{|f'(H)|^q,|f'(a)|^q\right\}\right]^{\frac{1}{q}}\right. \\
&\left.+\frac{b^2H^2}{b^{2p}}C_3^{1/p}\left(1,1,p,\frac{2a}{a+b}\right)\left[\sup\left\{|f'(H)|^q,|f'(b)|^q\right\}\right]^{\frac{1}{q}}\right\}.
\end{aligned}$$

Corollary 11. *Let the assumptions of Theorem 6 hold. If $|f'(x)| \leq M$ for all $x \in [a,b]$ and $\lambda = 0$, then we get the following Ostrowski type inequality for fractional from the inequality (2.7) integrals*

$$\begin{aligned}
&\left|\left[\left(\frac{x-a}{ax}\right)^{\alpha} + \left(\frac{b-x}{bx}\right)^{\alpha}\right]f(x) - \left(\frac{ab}{b-a}\right)^{\alpha} 2^{\alpha-1}\Gamma(\alpha+1)\left[J^{\alpha}_{1/x+}(f\circ g)(1/a) + J^{\alpha}_{1/x-}(f\circ g)(1/b)\right]\right| \\
\leq\ & \frac{M}{(\alpha+1)^{\frac{1}{q}}}\left[\frac{(x-a)^{\alpha+1}}{(ax)^{\alpha-1}x^{2p}}C_2^{1/p}\left(\alpha,0,p,\frac{a}{x}\right) + \frac{(b-x)^{\alpha+1}}{(bx)^{\alpha-1}b^{2p}}C_3^{1/p}\left(\alpha,0,p,\frac{x}{b}\right)\right].
\end{aligned}$$

References

[1] M. Alomaria, M. Darus, S.S. Dragomir, P. Cerone, Ostrowski type inequalities for functions whose derivatives are s-convex in the second sense, Applied Mathematics Letters 23 (2010) 1071–1076.

[2] M.W. Alomari, M.Darus, U.S. Kirmaci, Refinements of Hadamard-type inequalities for quasi-convex functions with applications to trapezoidal formula and to special means, Computers and Mathematics with Applications, 59 (2010) 225-232.

[3] M. Alomari, S. Hussain, Two inequalities of Simpson type for quasi-convex functions and applications, Applied Mathematics E-Notes, 11(2011), 110-117.

[4] D.A. Ion, Some estimates onthe Hermite-Hadamard inequality through quasi-convex functions, Annals of University of Craiova, Math. Comp. Sci. Ser. 34 (2007) 82-87.

[5] İ. İşcan, Hermite-Hadamard type inequalities for harmonically convex functions, Hacet. J. Math. Stat., Accepted for publication, 2013a.

[6] İ. İşcan, On generalization of different type inequalities for harmonically convex functions via fractional integrals, preprint 2013.

[7] İ. İşcan, Generalization of different type integral inequalities for s-convex functions via fractional integrals, Applicable Analysis. DOI:10.1080/00036811.2013.851785. Available online at: http://dx.doi.org/10.1080/00036811.2013.851785.

[8] İ. İşcan, New general integral inequalities for quasi-geometrically convex functions via fractional integrals, Journal of Inequalities and Applications, 2013, 2013:491, doi:10.1186/1029-242X-2013-491.

[9] İ. İşcan, On generalization of some integral inequalities for quasi-convex functions and their applications, International Journal of Engineering and Applied sciences (EAAS), 3 (1) (2013), 37-42.
[10] İ. İşcan, Generalization of different type integral inequalities via fractional integrals for functions whose second derivatives absolute values are quasi-convex, Konuralp journal of Mathematics, Volume 1 No. 2 pp. 67–79 (2013)
[11] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, Theory and applications of fractional differential equations, Amsterdam, Elsevier, 2006.
[12] M.Z. Sarıkaya and N. Aktan, On the generalization of some integral inequalities and their applications, Mathematical and Computer Modelling, 54 (2011) 2175- 2182.
[13] E. Set, New inequalities of Ostrowski type for mapping whose derivatives are s-convex in the second sense via fractional integrals, Computers and Math. with Appl. 63 (2012) 1147-1154.
[14] M.Z. Sarıkaya and H. Ogunmez, On new inequalities via Riemann-Liouville fractional integration, Abstract an Applied Analysis, vol. 2012, Article ID 428983, 10 pages, doi:10.1155/2012/428983.
[15] E. Set, M.E. Ozdemir and M.Z. Sarıkaya, On new inequalities of Simpson's type for quasi-convex functions with applications, Tamkang Journal of Mathematics, 43 (3) (2012) 357-364.
[16] M.Z. Sarıkaya, E. Set, H. Yaldız and N. Başak, Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities, Math. Comput. Model. 57(9-10) (2013) 2403-2407
[17] T.-Y. Zhang, , A.-P. Ji, and F. Qi, Integral inequalities of Hermite-Hadamard type for harmonically quasi-convex functions, Proc. Jangjeon Math. Soc. Vol. 16. No. 3. (2013) 399-407.

Department of Mathematics, Faculty of Sciences and Arts, Giresun University, Giresun, Turkey
E-mail address: imdat.iscan@giresun.edu.tr